



PLANT HEALTH INSECT CONTROL – CORN (FIELD, POP, SWEET, & SEED)

DUAL-ACTION, BROAD-SPECTRUM, LONG-RESIDUAL, FULL-SEASON YIELD PROTECTION

Insecticide + Fungicide Comparison



On-Farm Grower Trials



Application Recommendations

- 4.0-6.1 oz./A** Cutworms, flea beetles, Western bean cutworms, grasshoppers and stalk borers
- 4.0-10.3 oz./A** Aphids, armyworms, corn rootworms (adults), cereal leaf beetles, chinch bugs, European corn borers, corn earworms, hornworms, Japanese beetles (adults), Southern corn leaf beetles, corn blotch leafminer (adult), sap beetles, Southwestern corn borers, stink bugs and webworms
- 10.3 oz./A** Banks grass mites, carmine mites, two-spotted spider mites, lygus species.
Tank mix with **Dimethoate 400 EC** at 1 pt./A for two modes of action, improved control and resistance management

12 hour REI; 30-day PHI grain/stover; 60-day PHI forage/silage; 3-day PHI sweet corn.

Special Information

Apply using sufficient carrier to obtain thorough coverage of foliage. Minimum of 10, up to 20 gals./A by ground or 2, up to 5 gals./A by air. May include COC or equivalent at 1-2 qts. Can be tank mixed with commonly used herbicides, fungicides and foliar fertilizers. Do not apply more than 10.3 oz./A per application. Do not apply more than 41.2 oz./A per season including at-plant applications. 27.4 oz./A max seasonal for sweet corn, foliar use only.

Refer to the label maximum usage tables when using more than one product containing zeta cypermethrin or bifenthrin on this crop



2016 Hero Insecticide Early Corn Pest Preventative Assurance Program: See actual program for complete details.

Hero Insecticide is a Restricted Use Pesticide.

Always read and follow label directions. Hero Insecticide is not registered for use in California. FMC and Hero are trademarks of FMC Corporation or an affiliate. Headline is a trademark of BASF Corporation. ©2017 FMC Corporation. All rights reserved.