



INSECT CONTROL – CORN (FIELD, POP, SWEET & SEED)

BROAD-SPECTRUM, ECONOMICAL, EARLY & LATE SEASON INSECT CONTROL WITH SHORT PHI



Application Recommendations

1.28-2.8 oz./A	Cutworms: preventative or rescue.
1.8-4.0 oz./A	Corn earworms, green cloverworms, Western bean cutworms.
2.7-4.0 oz./A	Cereal leaf beetles, European corn borers, Southwestern corn borers, corn rootworms (adult), flea beetles, grasshoppers, hop vine borers, hornworms, Japanese beetles (adult), sap beetles (adult), Southern corn leaf beetles, stalk borers, stink bugs, tobacco budworms, webworms, aphids.
3.2-4.0 oz./A	Armyworms, chinch bugs.
	12-hour re-entry interval
	7-day PHI field and seed corn grain/stover/forage. Especially useful for late season seed corn protection. 3-day PHI sweet corn.
Special Information	Do not apply more than 4 oz./A per application or 16 oz./A per season on field, pop or corn grown for seed. Do not apply more than 4 oz./A per application or 24 oz./A per season on sweet corn.
	Apply by air or ground equipment using sufficient water to obtain thorough coverage of foliage. Minimum of 10, up to 20 gal./A by ground, or minimum of 2, up to 5 gal./A by air). May be tank mixed with commonly used herbicides, fungicides and foliar fertilizers. Chemigation allowed. See label requirements.



Mustang Maxx insecticide is a Restricted Use Pesticide.

Always read and follow label directions. Mustang Maxx insecticide is not registered for use in California. FMC and Mustang are trademarks of FMC Corporation or an affiliate. ©2017 FMC Corporation. All rights reserved.

