




INSECT CONTROL – SOYBEAN

BROAD-SPECTRUM, ECONOMICAL, EARLY AND LATE SEASON INSECT CONTROL

Stink Bug Control



On-Farm Grower Trials

| Treatment | Yield Bu/A |
|---|------------|
| Standard Program Delayed burndown fb glyphosate | 63.6 |
| Intensively Managed Authority® First DF Herbicide fb post glyphosate + Mustang® Maxx Insecticide + Headline® Fungicide | 68.7 |



Untreated vs. Mustang® Maxx Insecticide - 26 DAT


Application Recommendations

1.28-4.0 oz/A Cutworm, thistle caterpillar, silverspotted skipper.

2.8-4.0 oz/A Alfalfa caterpillar, southern armyworm, true armyworm, yellowstriped armyworm, bean leaf beetle, blister beetle, Colorado potato beetle, cucumber beetle, European corn borer, corn earworm, corn rootworm beetle (adult), flea beetle, green cloverworm, hornworms, Japanese beetle, leafhopper, leafminers (adults), Mexican bean beetle, potato leafhopper, seedcorn maggot (adult), soybean aphid, tobacco budworm, three-cornered alfalfa hopper, webworm, woolly bear caterpillar.

3.2-4.0 oz/A Beet armyworm, fall armyworm, grasshopper, looper, stink bugs, thrips, whitefly.

Special Information
12 hour REI, 21 day PHI.
 Apply using sufficient carrier to obtain thorough coverage of foliage to achieve control, minimum of 10 gals/A by ground or 2 gals/A by air (include COC or equivalent). Can be tank mixed with commonly used herbicides, fungicides, and foliar fertilizers. Do not make applications less than 7 days apart. Do not graze or harvest treated soybean forage, straw, or hay for livestock feed. Do not apply more than 24 oz/A per season.



Karate and Mustang Maxx are restricted use pesticides.

Always read and follow label directions. FMC, Authority and Mustang are trademarks of FMC Corporation or an affiliate. Headline is a trademark of BASF Corporation. Karate is a trademark of Syngenta Group Company. Orthene is a trademark of OMS Investments Inc. ©2017 FMC Corporation. All rights reserved.

