



Topguard® EQ Fungicide Technical Data

Corn



Topguard EQ Fungicide Technical Data

Corn




Target Diseases:

Common Rust
Southern Rust
Anthracnose Leaf Blight
Gray Leaf Spot
Northern Corn Leaf Blight
Northern Corn Leaf Spot
Southern Corn Leaf Blight
Eye Spot



Topguard EQ Technical Data

Corn



Application Guidelines:

Pre-harvest Interval: Seven days

Re-entry Interval: 12 hours (five days for detasseling seed corn)

Product use rates: 5-7 fl. oz./A

Restrictions:

Do not exceed 7 fl. oz./A for a single application.

Do not apply more than two applications of product per growing season.

Do not apply more than 14 fl. oz./A/year.

Please see product label for a complete list of restrictions.

Label:

Active Ingredient: Flutriafol and azoxystrobin

Mode of Action: Strobilurin inhibits fungal cell respiration. Triazole inhibits ergosterol production in cell membranes.

FRAC Group: 3 (Flutriafol); 11 (Azoxystrobin)

Learn more about Topguard EQ fungicide.

Contact your FMC Star Retailer, call 888-59-FMC-AG or visit FMCcrop.com.

TOPGUARD® EQ
FUNGICIDE

Always read and follow label directions. Topguard EQ fungicide is not registered for sale or use in California. FMC and Topguard are trademarks and Investing in farming's future is a service mark of FMC Corporation or an affiliate. Fortix is a trademark of Arysta LifeScience. Headline is a registered trademark of BASF. ©2016 FMC Corporation. All rights reserved. 17-FMC-1344 12/16

Investing in farming's future.™

