


Topguard® EQ Fungicide Technical Data

Pecan


Pecan Nut Scab Severity Hahira, GA



Topguard EQ Fungicide Technical Data

Pecan

Pecan, Scab Severity 2015 Tim Brennenman, UGA



Label contains fl. oz./A rate. Luna Sensation fungicide included NIS.
All treatments full season alternated with a Super Tin® + Elast® tank mix.
LSD (0.05)

Target Diseases:


Pecan Scab
Powdery Mildew
Anthracnose
Downy Spot
Zonate Leaf Spot



Topguard EQ Fungicide Technical Data

Pecan

Pecan Leaf Scab Severity Hahira, GA



Application Guidelines:

Pre-harvest Interval: 45 days

Re-entry Interval: 12 hours

Product use rates: 5-8 fl. oz./A

Restrictions:

No single application may exceed 8 fl. oz./A year

Do not apply more than 0.456 lb. a.i. of flutriafol/A/ year.

Do not apply more than 1.2 lbs. a.i. of azoxystrobin/A/ year.

Please see product label for a complete list of restrictions.

Label:

Active Ingredient: Flutriafol and azoxystrobin

Mode of Action: Strobilurin inhibits fungal cell respiration. Triazole inhibits ergosterol production in cell membranes.

FRAC Group: 3 (Flutriafol); 11 (Azoxystrobin)

Learn more about Topguard EQ fungicide.

Contact your FMC Star Retailer, call 888-59-FMC-AG or visit FMCcrop.com.

TOPGUARD® EQ
FUNGICIDE

Always read and follow label directions. Topguard EQ fungicide is not registered for sale or use in California. FMC and Topguard are trademarks and Investing in farming's future is a service mark of FMC Corporation or an affiliate. Quadris Top is a trademark of Syngenta Group Company. Luna Sensation is a trademark of Bayer. SUPER TIN is a trademark of United Phosphorus, Inc. ELAST is a trademark of Aceto Agricultural Chemicals Corporation. ©2016 FMC Corporation. All rights reserved. 17-FMC-1344 12/16

Investing in farming's future.™

FMC