


Topguard® EQ Fungicide Technical Data

Soybeans


Cercospora Severity
2014 Costello, Pioneer, LA



Topguard EQ Technical Data

Soybeans

Cercospora Severity
2014 Buckeye, Troy, OH



Target Diseases:

Aerial Blight
Alternaria Leaf Spot
Anthracnose
Brown Spot
Cercospora Blight
Purple Seed Stain
Frogeye Leaf Spot
Pod and Stem Blight
Powdery Mildew
Target Spot
Rust


Suppression:

White Mold/Stem Rot
Southern Blight

Topguard EQ Fungicide Technical Data

Soybeans

Cercospora Trial
2014 Costello, Pioneer, LA



Application Guidelines:

Pre-harvest Interval: 21 days

Re-entry Interval: 12 hours

Product use rates: 5-7 fl. oz./A

Restrictions:

Do not feed forage or hay to animals or permit animals to graze.

Do not apply more than 15.5 fl. oz./A/year.

Only one application of product at 8 fl. oz./A may be made to any one field during a single growing season.

Please see product label for a complete list of restrictions.

Label:

Active Ingredient: Flutriafol and azoxystrobin

Mode of Action: Strobilurin inhibits fungal cell respiration. Triazole inhibits ergosterol production in cell membranes.

FRAC Group: 3 (Flutriafol); 11 (Azoxystrobin)

Learn more about Topguard EQ fungicide.

Contact your FMC Star Retailer, call 888-59-FMC-AG or visit FMCcrop.com.

TOPGUARD® EQ
FUNGICIDE

Always read and follow label directions. Topguard EQ fungicide is not registered for sale or use in California. FMC and Topguard are trademarks and Investing in farming's future is a service mark of FMC Corporation or an affiliate. Fortix is a trademark of Arysta LifeScience. Quadris Top is a trademark of Syngenta Group Company. ©2016 FMC Corporation. All rights reserved. 17-FMC-1344 12/16

Investing in farming's future.™

FMC